

**NOVEL USE OF MIFEPRISTONE AND DERIVATIVES THEREOF AS
HEDGEHOG PROTEIN SIGNALING PATHWAY MODULATORS AND
APPLICATIONS OF SAME**

5 The present invention relates to a novel use of
mifepristone and derivatives thereof as Hedgehog
protein signaling pathway modulators, and also to
applications thereof for the treatment of pathologies
involving a tissue dysfunction associated with
10 deregulation of this pathway.

The Hedgehog protein signaling pathway plays an
essential role in the development of many embryonic
tissues, in particular in the nervous system, and is
15 thought to be involved in tissue maintenance and repair
in adults.

The Hedgehog gene (*Hh*), which was isolated in
drosophila and then in vertebrates, encodes a family of
20 secreted and very conserved signaling proteins, the
Hedgehog (*Hh*) proteins, which act as morphogens in many
tissues (for a review, see: Ingham et al., *Genes Dev.*,
2001, 15, 3059-3087; Marti et al., *Trends Neurosci.*,
2002, 25, 89-96; Weschler et al., *Annu Rev Neurosci.*,
25 2001, 24, 385-428). The *Hh* proteins are synthesized in
the form of immature precursors of approximately 45 kDa
that are autoprocessed into an active amino-terminal
form (*HhNp* for *N-terminal processed domain*), which is
doubly modified by the addition of two lipids: a
30 cholesterol residue (carboxy-terminal ester linkage)
and a palmitate (amino-terminal amide linkage). In
mammals, three families of *Hh* proteins exist: Sonic
(*Shh*), Indian (*Ihh*) and Desert (*Dhh*). *Shh* promotes
ventralization of the neural tube, specifying the early
35 phenotype of several types of neuron along the ventral
median line (spinal cord motoneurons, dopaminergic or
cholinergic neurons) and inducing the generation of
oligodendrocyte precursors from the ventral spinal
cord. Moreover, *Shh* induces gabaergic and dopaminergic

neuron survival, directs the outcome of serotonergic precursors and averts dopaminergic neuron death caused by the MPP toxin. Finally, it induces granular cell precursor proliferation in the early post-natal cerebellum. The other members of the Hedgehog family contribute, for their part, respectively to the development of bone tissue (Ihh), of the testes and of peripheral nerves (Dhh).

10 More recently, the Shh pathway has been identified in the adult brain, where the active amino-terminal form of the molecule is expressed in many regions of the mature nervous system, at a higher level than that encountered during the early post-natal period
15 (Traiffort et al., Eur. J. Neurosci.: 1999: 11, 3199-3214 and 2001, 14, 839-850). Although the roles of Shh in adults have not been completely elucidated, it first emerged, like other neurotrophic molecules, as a factor capable of promoting the survival and maintenance of
20 the phenotype of nervous system cells (Reilly et al., Mol. Cell. Neurosci., 2002, 19, 88-96; Charytoniuk et al., Eur. J. Neurosci., 2002, 16, 2351-2357). Under pathological conditions, such as a model of Parkinson's disease or a model of peripheral neuropathy, Shh is
25 capable of preserving the axonal projections of dopaminergic neurons in the striatum or of improving the time required for motor recovery subsequent to crush of the sciatic nerve (Tsuboi et al., Exp. Neurol., 2002, 173, 95-104; Pepinski et al., J. Pharm.
30 Sci., 2002, 91, 371-387).

The cellular response to the Hedgehog morphogen is controlled by the expression products of the *Patched* (*Ptc*) gene, which is a tumor suppressor gene, and of
35 the *Smoothed* (*Smo*) proto-oncogene; however, the exact mechanism of regulation of the Hedgehog pathway has not been completely elucidated. In mammals, two *Patched* genes exist, encoding, respectively, *Ptc1* and *Ptc2*, which are glycoproteins comprising 12 transmembrane

domains, homologous to bacterial transporters. The product of the *Smo* gene, which encodes a protein of the G protein-coupled receptor family, has no known endogenous ligand. In the absence of Hedgehog, Ptc is thought to block the constitutive activity of Smo. The binding of Hedgehog to Ptc is thought to lift this inhibition and allow signal transduction via Smo. The mechanism of regulation of Smo activity by Ptc, in mammals, could involve a molecule that is transported by Ptc and that interacts with Smo (Taipale et al., Nature, 2002, 418, 892-896). Activation of the Gli transcription factors is involved in the cascade of events resulting from the activity of Smo. The type I transmembrane protein HIP (Hedgehog-Interacting Protein) constitutes another receptor for Hedgehog molecules, which it binds with an affinity comparable to that of Ptc; HIP has been proposed as a negative regulator of the pathway (Ingham et al., mentioned above; Ho et al., Curr. Opin. Neurobiol., 2002, 12, 57-63; Taipale et al., Nature, 2001, 411, 349-354). In addition, the products of the *dispatched* (*disp*) gene, in particular DispA, are thought to be involved in the release and accumulation in the extracellular medium of Hedgehog proteins in soluble form (Ma et al., Cell, 2002, 111, 63-75).

Dysfunctions of the Shh signaling pathway have been linked to many cancers, in particular subsequent to the characterization of *Ptc* as a tumor suppressor gene. Specifically, inactivating mutations of *Ptc* are linked to Gorlin syndrome or basal cell nevus syndrome, an autosomal dominant disease characterized by craniofacial and cerebral malformations, but especially by a high incidence of various tumors, more particularly basal cell carcinomas of the skin and medulloblastomas in the brain. Mice heterozygotes for the *Ptc* gene develop cerebellar tumors, suggesting that a modification of the Shh pathway is the cause of these tumors (Goodrich et al., Science, 1997, 277, 1109-1113).

Mutations in the human *Ptc* or *Smo* genes are also observed in primitive neuroectodermal tumors of the central nervous system, mainly medulloblastomas (30% of cases), but also in sporadic forms of basal cell carcinomas (40% and 20%, respectively, for *Ptc* and *Smo*). In addition, *Shh* mutations (H133Y) are also linked to basal cell carcinomas. The *Smo* mutations that involve mainly two amino acids in the seventh hydrophobic domain of the receptor (W535L and S533N) induce constitutive activation of the pathway, which escapes the negative control by *Ptc*. On the other hand, *Ptc* mutations result in a decrease in the inhibition that it exerts on *Smo* in the absence of *Shh*. In the two cases, activation of the *Shh* pathway results therefrom and leads to a powerful mitogenic activity demonstrated in cultures of developing cerebellar granular cell precursors, and to blocking of the terminal differentiation step of these neuroblasts (Traiffort et al., Eur. J.; Neurosci., 1999, mentioned above; Charytoniuk et al., J. Physiol. Paris, 2002, 96, 9-16; Dahmane et al., development, 1999, 126, 3089-3100; Wallace et al., Curr. Biol., 1999, 22, 103-114; Weshler-Reya et al., Neuron., 1999, 22, 103-114). Similarly, the expression of *Smo* carrying one of these mutations in transgenic mice results in the presence of basal cell carcinomas, which demonstrates the direct involvement of *Smo* in the development of these tumors (Xie et al., Nature, 1998, 391, 90-92).

Besides basal cell carcinomas and medulloblastomas, other types of tumors have been linked to a deficiency in the Hedgehog signaling pathway; the location of these tumors is tightly correlated with the sites of expression of the components of the pathway during embryonic development. By way of nonlimiting example, mention may be made of: breast cancers and meningiomas related to *Ptc* mutations, and glioblastomas related to *Gli* mutations.

Because of the essential role of the Hedgehog protein signaling pathway in many physiological processes and, consequently, of the importance of the pathologies linked to a dysfunction thereof, the components of this pathway, such as the Smoothened or Patched (Patched 1 and Patched 2) proteins, the Dispatched (Dispatched 1 and Dispatched 2) proteins or else the HIP protein, represent targets for the development of novel molecules capable of modulating (activating or inhibiting) this pathway and therefore of positively or negatively regulating the development [proliferation, differentiation, migration, survival (apoptosis)] and/or the activity of differentiated cells and of stem cells, *in vitro* and/or *in vivo* in the embryo or in the adult.

Such molecules are useful in the treatment of tumors related to hyperactivation of the Hedgehog pathway: nervous tissue tumors (medulloblastomas, primitive neuroectodermal tumors, glioblastomas, meningiomas and oligodendrogliomas), skin tumors (basal cell carcinomas, trichoepitheliomas), muscle and bone tissue tumors (rhabdomyosarcomas, osteosarcomas) and tumors of other tissues (kidney, bladder).

Such molecules are also useful in the treatment of neurodegenerative-type pathologies requiring blocking of the Hedgehog pathway (Parkinson's disease, Huntington's chorea, Alzheimer's disease, multiple sclerosis, motoneuron disease), and diseases in which blocking of the Hedgehog signaling pathway could be beneficial, such as diabetes.

Such molecules are also useful in the medical or surgical treatment (plastic or healing surgery, tissue or organ transplantation) of many acute, subacute or chronic, genetic or acquired pathologies - involving a tissue dysfunction linked to deregulation of the Hedgehog pathway -, for inducing the formation, the

regeneration, the repair and/or an increase in the activity of tissues such as, without implied limitation: nervous tissue [central nervous system (brain) and peripheral nervous system (sensory, motor, sympathetic neurons)], bone, cartilage, testes, liver, spleen, intestine, pancreas, kidneys, smooth muscle and skeletal muscle, heart, lungs, skin and body hair system, mucous membranes, blood cells and cells of the immune system. By way of nonlimiting example of these pathologies, mention may in particular be made of neuropathies and related neuromuscular diseases, diabetes, alopecia, burns, ulcerations (skin and mucous membranes) and spermatogenesis disorders.

Various molecules capable of modulating the activity of the Hedgehog pathway have been identified:

- the Hedgehog proteins and derived polypeptides (fragments, variants, etc.), in particular Hedgehog protein agonists and antagonists (PCT international application WO 01/98344 in the name of Biogen); because of their size, these proteins and the derived polypeptides cannot cross the blood-brain barrier and cannot therefore be administered systemically, in particular for the treatment of brain tumors linked to hyperactivation of the Hedgehog protein signaling pathway. In addition, such molecules are difficult to produce and to purify and are relatively unstable;

- heterocyclic organic molecules (PCT international application WO 01/74344 in the name of Curis and Chen et al., PNAS, 2002, 99, 14071-14076);

- nitrogenous heterocyclic molecules (PCT international applications WO 01/19800, WO 01/26644 and WO 02/30421 in the name of Curis and Kamenetsky et al., J. Biol., 2002, 1, 1-19) and

- plant steroids derived from *Veratrum* spp (jervine,

cyclopamine and cycloposine) and from *Solanum* spp. (solanidine), substituted in the 16-, 17- or 18-position with an amine or an amine derivative, and cholesterol: American patent US 6,432,970 and PCT international applications WO 99/52534 and WO 01/27135 in the name of Johns Hopkins University School of Medicine; American patent US 6,291,516 and PCT international application WO 00/41545 in the name of Ontogeny and PCT international application WO 02/30462 in the name of Curis; Talpale et al., Nature, 2000, 406, 1005-1009; Berman et al., Science, 2002, 297, 1559-1561. However, cyclopamine is a teratogenic agent that causes holoprosencephaly and cyclopia in mammalian embryos, and an absence of toxicity, for mammals, has not been demonstrated for the other plant steroid-derived compounds.

It emerges from the above that no molecule currently exists that is effective for the treatment of pathological conditions requiring modulation of the activity of the Hedgehog protein signaling pathway, and for which a lack of toxicity has been verified by means of clinical trials in humans. Consequently, the inventors gave themselves the aim of providing compounds that better satisfy practical needs, in particular in that they are devoid of toxicity in humans and can therefore potentially be used in human therapy.

European patent EP 0 057 115 and American patent US 4,386,085 in the name of Roussel-Uclaf describe 19-nor steroid compounds substituted in the 11 β -position with a radical comprising a nitrogen, phosphorus or silicon atom, having anti-glucocorticoid, progestomimetic or anti-progestomimetic, androgenic or anti-androgenic properties. Because of these properties, the use of these steroid compounds is recommended for: combating conditions linked to hypersecretion of corticosteroids, especially for combating aging, in particular combating hypertension, atherosclerosis,

diabetes, obesity and also immunodepression and insomnia, for preparing contraceptives, for combating hormone deregulation, especially hormone-dependent cancers, amenorrhea, dysmenorrhea and luteal
5 insufficiency, and also for treating hypertrophies and cancer of the prostate, hyperandrogenism, anemia, hirsutism and acne.


Among these compounds, mifepristone (17β -hydroxy- 11β -(4-dimethylaminophenyl)- 17α -(prop-1-ynyl)estra-4,9-dien-3-one), also called RU-486 or RU-38486, is a 19-nor steroid substituted in the 11β -position with a radical comprising a nitrogen atom, which has a progesterone receptor antagonist activity and also a strong affinity for
15 glucocorticoid receptors and a weak affinity for androgen receptors; this molecule has been given a marketing authorization for termination of pregnancy, and phase II clinical trials are ongoing for the treatment, in menopausal women, of metastatic breast cancers.

20 The inventors have shown that mifepristone and some of its derivatives, that differ from the abovementioned cyclopamine-derived steroids in that they do not have a primary, secondary or tertiary amine function, or
25 nitrogenous heterocycle or amine derivatives substituted in the 16-, 17- and 18-position, are, surprisingly, active on the Hedgehog protein signaling pathway.

30 Because of its lack of toxicity in humans, mifepristone, which has been used in human therapy in various countries for many years, represents an advantageous alternative to cyclopamine and to its derivatives, for the treatment of pathological
35 conditions requiring a modulation (activation or inhibition) of the activity of the Hedgehog protein signaling pathway.

Consequently, a subject of the present invention is the


use of the compounds of formula (I)



in which:

- 5 - R_1 represents an organic radical containing from 1 to 18 carbon atoms, containing at least one nitrogen, phosphorus or silicon atom, the atom immediately adjacent to carbon 11 being a carbon atom,
- 10 - R_2 represents a hydrocarbon-based radical containing from 1 to 8 carbon atoms,


- X of formula:



15

represents the residue of a saturated or unsaturated, pentagonal or hexagonal ring optionally substituted with one or more groups chosen from the following radicals: C_1 - C_{12} alkyl, alkenyl, alkynyl, aryl or

20 aralkyl, hydroxyl $-OH$, carbonyl $\begin{array}{c} O \\ || \\ -C-H \end{array}$ or $\begin{array}{c} O \\ || \\ -C-X_1 \end{array}$, ether $-O-X_2$, ester $\begin{array}{c} O \\ || \\ -C-O-X_3 \end{array}$, oxycarbonyl $\begin{array}{c} O \\ || \\ O-C-X_4 \end{array}$, halogen, thiol, thioether $-S-X_5$, sulfinyl $\begin{array}{c} O \\ || \\ -S-X_6 \end{array}$, sulfonyl




, and also C₁-C₁₂ alkyl, alkenyl, alkynyl, aryl or aralkyl substituted with one or more hydroxyl -OH,

carbonyl $\begin{array}{c} \text{O} \\ \parallel \\ -\text{C}-\text{H} \end{array}$ or $\begin{array}{c} \text{O} \\ \parallel \\ -\text{C}-\text{X}_1 \end{array}$, ether -O-X₂, ester

$\begin{array}{c} \text{O} \\ \parallel \\ -\text{C}-\text{O}-\text{X}_3 \end{array}$, oxycarbonyl $\begin{array}{c} \text{O} \\ \parallel \\ \text{O}-\text{C}-\text{X}_4 \end{array}$, halogen, thiol,

5 thioether -S-X₅, sulfinyl $\begin{array}{c} \text{O} \\ \parallel \\ -\text{S}-\text{X}_{10} \end{array}$, or sulfonyl $\begin{array}{c} \text{O} \\ \parallel \\ -\text{S}-\text{X}_7 \\ \parallel \\ \text{O} \end{array}$.
functions, with X₁, X₂, X₃, X₄, X₅, X₆ and X₇ representing C₁-C₈ alkyl, C₂-C₈ alkenyl or C₂-C₈ alkynyl groups, or C₆-C₁₅ aryl or C₆-C₁₅ aralkyl groups,

10 - the group C=A in the 3-position represents an oxo group, which is free or blocked in the form of a ketal, an alcohol -CH-OH, ether -CH-O-Y₁, alkyl carboxylate



15 , C=NOH or C=NO-Y₃ group, or a CH₂ group, Y₁, Y₂ and Y₃ representing an alkyl radical containing from 1 to 8 carbon atoms or an aralkyl group containing from 7 to 15 carbon atoms, and

- B and C together form a double bond or an epoxide bridge, and of the derived salts, for preparing a
20 medicinal product capable of modulating (activating or inhibiting) the Hedgehog protein signaling pathway, intended for the treatment of pathologies involving a tissue dysfunction linked to deregulation of this pathway.

25

Among the derived salts, mention may be made of the addition salts with acids of the compounds of

formula I, for instance the salts formed with hydrochloric acid, hydrobromic acid, nitric acid, sulfuric acid, phosphoric acid, acetic acid, formic acid, propionic acid, benzoic acid, maleic acid, fumaric acid, succinic acid, tartaric acid, citric acid, oxalic acid, glyoxylic acid, aspartic acid, alkanesulfonic acids, such as methanesulfonic acid or ethanesulfonic acid, arylsulfonic acids, such as benzenesulfonic acid or para-toluenesulfonic acid, and arylcarboxylic acids.


According to an advantageous embodiment of the invention, R_2 represents a linear or branched, saturated alkyl radical containing from 1 to 4 carbon atoms, preferably a methyl radical.

According to another advantageous embodiment of the invention, X represents an optionally substituted pentagonal ring.

According to an advantageous arrangement of this embodiment, said pentagonal ring is substituted with at least one alkenyl or alkynyl group, preferably with an alkynyl group, preferably in the 17-position.

Advantageously, said pentagonal ring is also substituted with at least one hydroxyl group.


Preferably, X represents the residue of a pentagonal ring of formula:



According to yet another advantageous embodiment of the invention, R_1 represents a hydrocarbon-based radical containing from 1 to 18 carbon atoms and containing at least one nitrogen atom, selected from:

- the R₁ values which represent a primary, secondary or tertiary alkyl radical containing from 1 to 8 carbon atoms, in particular a methyl, ethyl, n-propyl, isopropyl, butyl, isobutyl, tert-butyl, pentyl, hexyl, cyclopropyl, cyclobutyl, cyclopentyl or cyclohexyl radical, containing at least one nitrogen atom or substituted with a heterocycle containing at least one nitrogen atom and optionally substituted with an alkyl radical containing from 1 to 8 carbon atoms, such as a methyl, ethyl or n-propyl radical, in particular 3,4- or 2-pyridyl radicals, the thiazolyl radical or the piperidiny radical, and

- the R₁ values which represent an aryl or aralkyl radical carrying an amine function, in particular a phenyl or benzyl radical carrying an amine function:




in which Z₁ and Z₂, which may be identical or different, represent a linear, branched or cyclic alkyl radical containing from 1 to 8 carbon atoms, it being possible for Z₁ and Z₂ to be optionally combined so as to form a heterocycle with the nitrogen atom. Preferably, Z₁ and Z₂ represent a C₁-C₄ alkyl radical, and preferentially a methyl radical.

According to yet another advantageous embodiment of the invention, the group C=A in the 3-position represents an oxo group.

According to yet another advantageous embodiment of the invention, B and C together form a double bond.

According to an advantageous arrangement of the above embodiments, said compound of formula (I) is 17β-hydroxy-11β-(4-dimethylaminophenyl)-17α-(prop-

1-ynyl)estra-4,9-dien-3-one of formula:



The compounds of formula I and their addition salts
5 with pharmaceutically acceptable acids are produced as
described in European patent 0 057 115 and US patent
4,386,085.

The compounds of formula I and their addition salts
10 with pharmaceutically acceptable acids which are
capable of modulating the activity of the Hedgehog
protein signaling pathway are used in the treatment of
tumors linked to hyperactivation of the Hedgehog path-
way, in particular nervous tissue tumors
15 (medulloblastomas, primitive neuroectodermal tumors,
glioblastomas, meningiomas and oligodendrogliomas),
skin tumors (basal cell carcinomas, tricho-
epitheliomas), muscle and bone tissue tumors
(rhabdomyosarcomas, osteosarcomas) and tumors of other
20 tissues (kidney, bladder).

Such compounds are also used in the treatment of
pathologies requiring blocking of the Hedgehog pathway,
in particular neurodegenerative-type pathologies such
25 as Parkinson's disease, Huntington's chorea,
Alzheimer's disease, multiple sclerosis and motoneuron
disease, or else other pathologies in which blocking of
the Hedgehog signaling pathway could be beneficial,
such as diabetes.

30

Such compounds are also used in the medical or surgical
treatment (plastic or healing surgery, tissue or organ
transplantation) of many acute, subacute or chronic,
genetic or acquired pathologies involving a tissue

dysfunction linked to deregulation of the Hedgehog pathway, for inducing, from stem cells or differentiated cells, the formation, the regeneration, the repair and/or an increase in the activity of tissues
5 such as, without implied limitation: nervous tissue [central nervous system (brain) and peripheral nervous system (sensory, motor, sympathetic neurons)], bone, cartilage, testes, liver, spleen, intestine, pancreas, kidneys, smooth muscle and skeletal muscle, heart,
10 lungs, skin and the body hair system, mucous membranes, blood cells and cells of the immune system. By way of nonlimiting example of these pathologies, mention may in particular be made of neuropathies and associated neuromuscular diseases, diabetes, alopecia, burns,
15 ulcerations (skin and mucous membranes) and spermatogenesis disorders.

The useful dosage varies according to the condition to be treated, to the route and rate of administration,
20 and also to the nature and the weight of the species to be treated (human or animal); it may, for example, vary from 100 mg to 1 g per day in adults, when administered orally.

25 The compounds of formula (I) and their salts are used via the digestive route (orally or sublingually), parenterally or locally. They may be in the form of simple or sugar-coated tablets, of gelatin capsules, of granules, of syrup, of suppositories, of injectable
30 preparations, of ointments, of creams, of gels or of an aerosol, which are prepared according to the usual methods.

In these pharmaceutical forms, the compounds of formula
35 (I) are incorporated into excipients conventionally used in pharmaceutical compositions, such as talc, gum arabic, lactose, starch, magnesium stearate, cocoa butter, aqueous or nonaqueous carriers, fatty substances of animal or plant origin, paraffin

derivatives, glycols, various wetting agents, dispersants or emulsifiers, or preserving agents.

5 The use of the compounds of formula (I) as defined above, and in particular of mifepristone, is more advantageous than that of the existing Hedgehog protein signaling pathway modulators in that these compounds are devoid of toxicity in humans and can therefore effectively be used in human therapy, unlike
10 cyclopamine and its derivatives, which are teratogenic.

Besides the above arrangements, the invention also comprises other arrangements which will emerge from the following description, which refers to examples of use
15 of mifepristone and of its derivatives according to the present invention, and also to the attached drawings in which:

- Figure 1 illustrates the inhibition, by
20 mifepristone, of the differentiation of mesenchymal cells (C3H10T1/2 line) induced by the Sonic Hedgehog (Shh) protein. Untransfected C3H10T1/2 cells (white columns) and those transfected with the Shh protein expression plasmid, called pRK5 (C3H10T1/2-SHH, grayish
25 columns), are treated for 5 days with mifepristone (RU 38 486, 10 μ M), cyclopamine (10 μ M) or their respective carriers, dimethyl sulfoxide (DMSO) or ethanol (EtOH), or are not treated. The differentiation of the C3H10T1/2 cells, which is accompanied by
30 induction of alkaline phosphatase activity, is then evaluated by measuring the optical density at 415 nm, in the presence of the alkaline phosphatase substrate, para-nitrophenyl phosphate. The values indicated correspond to the mean \pm standard error (n = 4), and

35

- Figure 2 illustrates the dose-effect curve for the inhibition of differentiation of the C3H10T1/2 cells by mifepristone. C3H10T1/2 cells transfected with the Shh protein expression plasmid, called pRK5, are treated

for 5 days with increasing amounts of mifepristone (RU 38 486, 1 nM to 10 μ M) or are not treated. The inhibition of differentiation of the C3H10T1/2 cells is expressed through the percentage alkaline phosphatase activity, relative to the nontreated cells (maximum activity = 100%). The values indicated correspond to the mean \pm standard error (n = 4) of two independent experiments. The concentration that inhibits 50% of cell differentiation (IC₅₀) is 3.7 μ M \pm 0.3 μ M.

EXAMPLE: EFFECT OF MIFEPRISTONE ON THE HEDGEHOG PROTEIN SIGNALING PATHWAY

The effect of mifepristone on the Hedgehog protein signaling pathway is determined by analyzing the differentiation of the C3H10T1/2 pluripotent fibroblast cell line, induced beforehand with the ShhN protein, in the presence or absence of mifepristone or of cyclopamine, for comparison.

1) Materials and methods

The C3H10T1/2 pluripotent fibroblast cell line (ATCC) is cultured under the conditions recommended by the ATCC. C3H10T1/2 cells were transfected with the plasmid pRK5 encoding the mouse Shh protein (Taipale et al., Nature, 2001, 411, 349-354), using calcium phosphate, according to conventional protocols. The cells were then cultured for 5 days in the presence of mifepristone (10 μ M), of cyclopamine (10 μ M) or else of corresponding volumes of their respective carriers, dimethyl sulfoxide (DMSO) and ethanol (EtOH); the alkaline phosphatase activity induced by the differentiation of the C3H10T1/2 cells into muscle, adipocyte, chondrocyte or osteoblast cells was then evaluated by measuring the optical density at 415 nm, in the presence of the alkaline phosphatase substrate, para-nitrophenyl phosphate. Alternatively, the dose-response curve for mifepristone was determined under the same experimental conditions, in the presence of

increasing concentrations of mifepristone (1 nM to 10 μ M).

2) Results

5 a) Comparative effect of mifepristone and of cyclopamine on C3H10T1/2 cell differentiation

The results given in figure 1 show that the 3- to 10-fold increase in alkaline phosphatase activity that is induced by the overexpression of Shh is completely
10 inhibited in the presence of 10 μ M of mifepristone or of cyclopamine.

These results demonstrate that mifepristone, which, unlike cyclopamine, is devoid of any teratogenic
15 effect, is capable, like cyclopamine, of blocking the Shh protein-induced differentiation of mesenchymal cells. Mifepristone therefore represents a Hedgehog protein signaling pathway antagonist that is devoid of any teratogenic effect and that is useful for the
20 treatment of pathologies requiring blocking of the Hedgehog pathway, such as cancer, neurodegenerative diseases and diabetes.

25 b) Dose-effect curve for the inhibition of C3H10T1/2 cell differentiation by mifepristone

The results given in figure 2, which confirm those given in figure 1, show that a concentration of 10 μ M of mifepristone completely inhibits the activation of alkaline phosphatase induced by the overexpression of
30 the Shh protein. The concentration that inhibits 50% of cell differentiation (IC_{50}) is 3.7 μ M \pm 0.3 μ M.

As emerges from the above, the invention is in no way limited to its methods of implementation, execution and
35 application which have just been described more explicitly; on the contrary, it encompasses all the variants thereof which may occur to those skilled in the art, without departing from the context or the scope of the present invention.